



**E Journal  
of Cardiovascular  
Medicine**

| Volume **5** | Issue **3** |

| July-September **2017** |

[www.ejcvsmmed.org](http://www.ejcvsmmed.org)

# **E Journal of Cardiovascular Medicine**

**Comparison between two sternum closing techniques  
by median sternotomy during open cardiac surgery**

*Kenan A. Kara1, Zafer Erk, Aytuğ Koçyiğit, Şenol Gülmen, Ahmet Öcal, Hüseyin Okutan*

**Hybrid treatment of aortic bifurcation aneurysm**

*Funda Yıldırım, Dilşad A. Şenarşlan, Adnan Taner Kurdal, Barış Bayram,  
Tülün Öztürk, Mustafa Cerrahoğlu, İhsan İşkesen, Ömer Tetik*

**Endovascular treatment of a ruptured abdominal aortic  
aneurysm in a patient with cardiogenic shock**

*Ali B. Budak, Atike T. Kunt, Orhan E. Günertem, Naim B. Tümer, Kanat Özışık, Serdar Günaydın*



**E Journal of Cardiovascular Medicine**  
is a global e-journal targeting articles on:

- clinical cardiology,
- interventional cardiology,
  - arrhythmia,
- cardiovascular surgery,
- vascular & endovascular surgery,
- vascular biology

## Editor-in-Chief

### Prof. Öztekin Oto

*FESC, FACC / President, Heart and Health Foundation of Turkey / Izmir / Turkey*

## Asistant Editors

### Prof. Ali Kutsal

*Sami Ulus Children Hospital Department of  
Cardiovascular Sugery / Ankara / Turkey*

### Prof. Erdem Silistreli

*Dokuz Eylül University, Department of Cardiovascular  
Sugery / Izmir / Turkey*

### Prof. Bektaş Battaloğlu

*İnönü University, Department of Cardiovascular Sugery  
Malatya / Turkey*

### Dr. Onur Saydam

*Karaman State Hospital Cardiovascular Surgery  
Karaman / Turkey*

### Dr. Emre Doğan

*Trabzon Ahi Evren Cardiovascular Surgery Hospital  
Trabzon / Turkey*

### Dr. Taylan Adademir

*Kartal Koşuyolu Resarch Hospital / Istanbul / Turkey*

## Owner

© TÜSAV Heart and Health Foundation of Turkey

## Administration Office

Şair Eşref Bulvarı, 1402 Sk. No. 2/2 Özbaş Apt.  
Alsancak - Izmir / Turkey  
Tel: + 90 232 464 19 63 / Fax: +90 232 464 24 70  
e-mail: info@oztekinoto.com | info@tksv.org

## Publishing Coordinator

### Hüseyin Kandemir

huseyin@medikalakademi.com.tr

## Publisher

Medikal Akademi Yayıncılık ve Prodüksiyon Tic. Ltd. Sti.  
Halaskargazi Cad. No: 172, D: 134 - Şişli / İstanbul  
Tel: +90 537 309 29 55, Faks: (0212) 233 90 61  
www.medikalakademi.com.tr/hizmetlerimiz

## International Scientific Advisory Board

### Prof. Harald Kaemmerer

*German Heart Centre / Munich / Germany*

### Prof. Marko Turina

*University Hospital of Zurich / Zurich / Switzerland*

### Prof. Frank W. Selke

*Chief of Cardiothoracic Surgery at Brown Medical School  
Rhode Island / USA*

### Prof. Joseph E. Bavaria

*Hospital of the University of Pennsylvania  
Philadelphia / USA*

### Prof. Fausto Pinto

*Director of Lisbon Cardiovascular Institute / Portugal  
& President of the European Society of Cardiology*

### Prof. Lazar Davidovic

*Belgrade Medical School Cardiovascular Surgery  
Belgrade / Serbia*

### Prof. Stephan Schueler

*Tyne Freeman Hospital, Department for Cardiothoracic  
Surgery Newcastle / United Kingdom*

### Prof. Piotr Kasprzak

*University Hospital Regensburg, Director of Vascular  
Surgery / Regensburg / Germany*

### Prof. Jose Luis Pomar

*Hospital Clinico de Barcelona, Department of  
Cardiovascular Sugery / Barcelona / Spain*

**Prof. Mohamed Moustafa Abdelaal**

*Kafrelsheikh University Hospital, Cardiothoracic surgery and  
General Director / Kafr El Sheikh / Egypt*

**Assoc. Prof. Barış Akça**

*Inonu University School of Medicine, Department of  
Cardiovascular Surgery / Malatya / Turkey*

**Dr. Rezan Aksoy**

*Siyami Ersek Training and Research Hospital,  
Cardiovascular Surgery / Istanbul / Turkey*

**Dr. Şafak Alpat**

*Birmingham Children's Hospital Pediatric Cardiovascular  
Surgery / Birmingham / UK*

**Dr. Mustafa Aldemir**

*Kocatepe University, Department of Cardiovascular  
Surgery / Afyon / Turkey*

**Dr. Elena Zapata-Arriaza**

*Hospital Universitario virgen del Rocío, Instituto de  
biomedicina de Sevilla, Vascular Medicine / Sevilla / Spain*

**Dr. Mehmet Atay**

*Bakırköy Sadi Konuk Research Hospital, Cardiovascular  
Surgery / Istanbul / Turkey*

**Assoc. Prof. Hakan Aydın**

*Sami Ulus in Ankara Training and Research Hospital,  
Cardiovascular Surgery / Ankara / Turkey*

**Assoc. Prof. Ahmet Çağrı Aykan**

*Ahi Evren University of Health Sciences, Thoracic and  
Cardiovascular Surgery / Trabzon / Turkey*

**Assoc. Prof. Vedat Bakuy**

*Bakırköy Sadi Konuk Training and Research Hospital,  
Cardiovascular Surgery / Istanbul / Turkey*

**Dr. Stefano Bartoli**

*ASL Roma2, Cardiovascular Surgery  
Rome / Italy*

**Assoc. Prof. Elif Börekçi**

*Bozok University Research and Application Hospital,  
Internal Medicine / Yozgat / Turkey*

**Dr. Tomasa Centella**

*Hospital Ramón y Cajal, Cardiovascular Surgery  
Madrid / Spain*

**Assoc. Prof. Ahmet Çalışkan**

*Dicle University School of Medicine, Cardiovascular Surgery  
Diyarbakır / Turkey*

**Dr. Gökhan Çavuşoğlu**

*Ahi Evren University of Health Sciences, Thoracic and  
Cardiovascular Surgery, Radiology / Trabzon / Turkey*

**Dr. Deniz Çevirme**

*Kartal Koşuyolu Research and Education  
Hospital, Cardiovascular Surgery / Istanbul / Turkey*

**Prof. Ferit Çiçekçioğlu**

*Bozok University, Training and Research Hospital,  
Cardiovascular Sugery / Yozgat / Turkey*

**Assoc. Prof. Ertan Demirdaş**

*Bozok University Research and Application Hospital and  
Cardiovascular Surgery / Yozgat / Turkey*

**Assoc. Prof. Yüksel Dereli**

*Necmettin Erbakan University, Meram Midical Faculty  
Hospital, Cardiovascular Surgery / Konya / Turkey*

**Assist.Prof. İnci Selin Doğan**

*Karadeniz Technical University Faculty of Pharmacy  
Pharmacology, Medicinal Chemistry / Trabzon / Turkey*

**Dr. Vehbi Doğan**

*Sami Ulus Training and Research Hospital, Pediatric  
Cardiology / Ankara / Turkey*

**Dr. Çağrı Düzyol**

*Kocaeli Derince Education and Research Hospital  
Cardiovascular Surgery / Kocaeli / Turkey*

**Assoc. Prof. Hüseyin Ede**

*Bozok University, Medical Faculty, Cardiovascular Surgery  
Yozgat / Turkey*

**Dr. İlker Ertuğrul**

*Sami Ulus Training and Research Hospital, Pediatric  
Cardiology / Ankara / Turkey*

**Prof. Niyazi Görmüş**

*Necmettin Erbakan University, Meram Medical Faculty  
Hospital, Cardiovascular Surgery / Konya / Turkey*

**Assist. Prof. Adem Güler**

*Gulhane Military Medical Academy Department of  
Cardiovascular Surgery / Ankara / Turkey*

**Assoc. Prof. Mustafa Gülgün**

*GATA Department of Pediatrics, Division of Pediatric  
Cardiology / Ankara / Turkey*

**Prof. Usama Ali M. Hamza**

*Mansoura University Faculty of Medicine, Cardiothoracic  
Surgery, Cardiovascular Surgery / Mansoura / Egypt /*

**Dr. James B Hermiller**

*St Vincent's Medical Group, Interventional Cardiology  
Indianapolis / USA*

**Dr. Akihiko Ikeda**

*Tsukuba Medical Center Hospital, Cardiovascular  
Surgery / Tsukuba / Japan*

**Dr. Richard W Issitt**

*Great Ormond Street Hospital, Cardiac Surgery -  
Pediatric Cardiology / London / UK*

**Dr. Mehmet Kalender**

*Derince Training and Research Hospital, Cardiovascular  
Surgery / Kocaeli / Turkey*

**Dr. Ayşegül Karadeniz**

*Ahi Evren University of Health Sciences, Thoracic and  
Cardiovascular Surgery, Radiobiology / Trabzon / Turkey*

**Assoc. Prof. Osman Kayapınar**

*Duzce University, Medical Faculty Department of  
Cardiology / Düzce / Turkey*

**Assoc. Prof. Alper Kepez**

*Marmara University Training and Research Hospital Cardiol-  
ogy Clinic / Istanbul / Turkey*

**Assoc. Prof. Yasir Khan Khan**

*Ch. Pervaiz Elahi Institute of Cardiology, Cardiovascular Sur-  
gery / Punjab / Pakistan*

**Assoc. Prof. Levent Korkmaz**

*Ahi Evren University of Health Sciences, Thoracic and  
Cardiovascular Surgery / Trabzon / Turkey*

**Assoc. Prof. Ulaş Kumbasar**

*Hacettepe University Medical School Cardiovascular  
Surgery / Ankara / Turkey*

**Dr. Redha Lakehal**

*Department of heart surgery, EHS Erriadh / Constantine  
Algeria*

**Dr. Wei-Chieh Lee**

*Kaohsiung Chang Gung Memorial Hospital, Cardiology  
Kaohsiung City / Tayvan*

**Dr. José Luis Serrano Martínez**

*University Hospital of Granada, Department of Internal  
Medicine / Granada / Spain*

**Assoc. Prof. Ümit Mentşe**

*Ahi Evren University of Health Sciences, Thoracic and  
Cardiovascular Surgery / Trabzon / Turkey*

**Dr. Nooredin Mohammadi**

*Iran University of Medical Sciences, Cardiology, Demand  
for Health Care, Determinants of Health / Tehran / Iran*

**Dr. Vinod Namana**

*Maimonides Medical Center, Department of Medical  
Research / New York / USA*

**Dr. Silvio Nocco**

*Sirai Hospital, Department of Cardiology / Carbonia / Italy*

**Assoc. Prof. Zeynep Tuğba Özdemir**

*Bozok University School of Medicine, Internal Medicine  
Yozgat / Turkey*

**Dr. Tanıl Özer**

*Kartal Koşuyolu Yüksek İhtisas Research and Education  
Hospital / Istanbul / Turkey*

**Prof. Murat Özeren**

*Mersin University Medical School, Cardiovascular  
Surgery / Mersin / Turkey*

**Assoc. Prof. Emre Özker**

*Başkent University School of Medicine, Department of  
Cardiovascular Surgery / Ankara / Turkey*

**Dr. Abdullah Özyurt**

*Mersin Maternity and Children Diseases Hospital, Pediatric  
Cardiology / Mersin / Turkey*

**Dr. Recep Oktay Peker**

*Hacettepe University, Department of Cardiovascular Surgery  
Ankara / Turkey*

**Dr. Hikmet Sahratov**

*Gülhane Education and Research Hospital, Department of  
Cardiovascular Surgery / Ankara / Turkey*

**Dr. Gonzalo Luis Alonso Salinas**

*Marcelo Sanmartín of Hospital Universitario Ramón y Cajal  
Madrid / Spain*

**Dr. Stefano Salizzoni**

*Città della Salute e della Scienza, Cardiac Surgery,  
Cardiac Surgery / Turin / Italy*

**Dr. Gökhan Sargın**

*Adnan Menderes University Medical School, Internal  
Medicine / Aydın / Turkey*

**Dr. Mustafa Seren**

*Ankara 29 Mayıs State Hospital and Cardiovascular  
Surgery / Ankara / Turkey*

**Prof. Erdem Silistreli**

*Dokuz Eylül University, Department of Cardiovascular  
Surgery / İzmir / Turkey*

**Assoc. Prof. Ömer Tanyeli**

*Necmettin Erbakan University, Meram Medical Faculty  
Hospital, Cardiovascular Surgery / Konya / Turkey*

**Dr. İlker Tekin**

*Antalya Medicalpark Hospital, Cardiovascular Surgery  
Antalya / Turkey*

**Assist. Prof. Dinçer Uysal**

*Isparta Süleyman Demirel University, Department of  
Cardiovascular Surgery / Isparta / Turkey*

**Dr. Olivier Villemain**

*IM3C Necker-Enfants Malades, AP-HP, Université Paris  
Descartes, Pediatric Cardiology, Radiology / Paris / France*

**Dr. Mustafa Esat Yamaç**

*Ahi Evren University of Health Sciences, Thoracic and  
Cardiovascular Surgery / Trabzon / Turkey*

**Assoc. Prof. Ali Ümit Yener**

*Canakkale Onsekiz Mart University Medical Faculty,  
Department of Cardiovascular Surgery / Çanakkale / Turkey*

**Dr. Dilek Yeşilbursa**

*Uludağ University, Medical Faculty, Department of  
Cardiology / Bursa / Turkey*

**Dr. Mustafa Yılmaz**

*Sami Ulus Training and Research Hospital, Pediatric  
Cardiology / Ankara / Turkey*

| Volume **5** | Number **3** | July-September **2017** |

### Research Articles

**Comparison between two sternum closing techniques by median sternotomy during open cardiac surgery | 44**

Kenan A. Kara, Zafer Erk, Aytağ Koçyiğit, Şenol Gülmen, Ahmet Öcal, Hüseyin Okutan

### Case Report

**Hybrid treatment of aortic bifurcation aneurysm | 50**

Funda Yıldırım, Dilşad Amanvermez Şenarslan, Adnan Taner Kurdal,  
Barış Bayram, Tülün Öztürk, Mustafa Cerrahoğlu, İhsan İşkesen, Ömer Tetik

**Extra-anatomic left carotico-subclavian artery bypass technique in a patient with subclavian steal syndrome: Case report | 54**

Ertan Demirdaş, Kıvanç Atılğan, Zafer Cengiz Er, Ferit Çiçekçioğlu

**Use of new oral anticoagulants secondary to paget–schroetter syndrome in a young sportsman: A case report | 58**

Kenan A. Kara, Uğur Diliçikık, Başar Sarıkaya, Şenol Gülmen

**Endovascular treatment of a ruptured abdominal aortic aneurysm in a patient with cardiogenic shock | 62**

Ali Baran Budak, Atike Tekeli Kunt, Orhan Eren Günertem,  
Naim Boran Tümer, Kanat Özışık, Serdar Günaydın

# Comparison between two sternum closing techniques by median sternotomy during open cardiac surgery

Kenan Abdurrahman Kara<sup>1</sup>, Zafer Erk<sup>2</sup>, Aytuğ Koçyiğit<sup>3</sup>, Şenol Gülmen<sup>4</sup>, Ahmet Öcal<sup>1</sup>, Hüseyin Okutan<sup>5</sup>

<sup>1</sup>) Hisar Intercontinental Hospital, Department of Cardiovascular surgery, M.D., Ümraniye, Istanbul, Turkey

<sup>2</sup>) Antalya Research and Training Hospital, Department of Cardiovascular surgery, M.D., Antalya, Turkey

<sup>3</sup>) Tekden Hospital, Department of Cardiovascular surgery, M.D., Denizli, Turkey

<sup>4</sup>) Süleyman Demirel University Hospital, Department of Cardiovascular surgery, M.D., Isparta, Turkey

<sup>5</sup>) Medicalpark Hospital, Department of Cardiovascular surgery, M.D., Antalya, Turkey

## Abstract

**Introduction:** Recovery failures such as sternal dehiscence, sternal pain, sternal wound infections, osteomyelitis and mediastinitis can be seen after median sternotomy in coronary bypass surgery. In this study, two sternum band closure methods which we find more effective and secure than conventional sternal closure method, are compared.

**Method:** In this prospective study that compares two sternal band closure techniques, 114 patients are divided in two groups as Group A (n=54) and Group B (n=60). This research compares the results for sternal dehiscence, sternal wound infections, sternal pain and mediastinitis in these patient groups between December 2011 and July 2012.

**Results:** Out of the two, a meaningful recovery is achieved in Group A compared to Group B (p<0.05). No superficial wound infection, sternal dehiscence and mediastinitis are found in patients of both band groups.

**Conclusion:** In this study, two different band closure techniques for sternotomy closure after the coronary bypass surgery are compared, and it is revealed that sternal pain index decreases meaningfully, compared to the other.

**Key Words:** Sternal band, sternal dehiscence, sternotomy, pain index

Kara K.A., Erk Z., Koçyiğit A., Gülmen Ş., Öcal A., Okutan H. Comparison between two sternum closing techniques by median sternotomy during open cardiac surgery EJCM 2017; 05 (3): 44-49. Doi: 10.15511/ejcm.17.00344.



## Introduction

Median sternotomy is the most frequently used incision in patients of open cardiac surgery. Following open cardiac surgery by median sternotomy, sternum is closed by surgical steel wires as whole layer. Potential post-operative sternal instability definition is dehiscence of both sternal parts due to cutting of sternal cortical layers by sternum closing wires with inappropriate movements of patient. This may result in conditions ranging from increased postoperative pain, sternal wound infection and mediastinitis associated with higher morbidity and mortality, prolonged hospital stay and increased treatment cost.<sup>(1-4)</sup>

Sternal bands used in our study are reported to prevent cutting of sternal cortex as tension on sternum is applied on larger and more balanced area, unlike to conventional sternal steel wires.<sup>(4-6)</sup>

The aim of this study to compare two different sternal band closing methods combined by conventional sternal closing method and to report the results.

## Patients and Method

In our clinic, 114 patients (60 males, 54 females; mean age 65,095 years) underwent to open cardiac surgery between December 2011 and July -2012 were prospectively reviewed. Sternal closing was performed by Peninsula Sterna –Band® (L.L.C. Livonia MI. USA ) was used in 54 patients consisting first group (group A) and by Ethicon Sternum Band (Norderstedt, Germany) used in 60 patients consisting second group (group B).

Patients with pre-operative renal insufficiency, left ventricle dysfunction, patients receiving anticoagulants, those who have hemorrhagic disorder and those who received postoperative over two drains, were not included in the study. Preoperative demographic properties of patients are shown in **Table 1**.

Anesthetic medication and prophylaxis were identical in both groups. Following standard sterilization, operating area was covered by ioban 2 drape (3M Health Care, St Paul, MN, USA). Following median sternotomy, antegrade cold blood cardioplegia with intervals of

**Table 1.** Preoperative demographic properties.

Group	A	B	p value
Patient number	54	60	NS
Age (Years)	64.96 ± 9.88	65.23 ± 10.11	NS
Weight (kg)	75.16 ± 12.	78.1 ± 17.48	NS
Height (cm)	165.00 ± 8.4	166.45 ± 9.25	NS
BMI	27.59 ± 3.98	28.12 ± 5.35	NS
COPD	21 (% 38.9)	28 ( % 46.7)	NS
DM	20 (% 37.0)	27 ( % 45)	NS
Obesity ( BMI > 30)	25 (% 46.3 )	23 ( % 38.3)	NS
Osteoporosis	3 (% 5.6 )	9 ( % 15 )	NS
Smaller Sternum	7 ( % 13 )	6 ( % 10)	NS

*COPD: Chronic obstructive pulmonary disease; DM: Diabetes mellitus; BMI: Body mass index; NS: Non-significant.*

20 minutes and continuous retrograde blood cardioplegia were used with standard cardiopulmonary bypass.

Retrosternal two sternal bands at intercostal spaces 3 and 4 and four standard 5/0 steel wires at intercostal spaces 1- 2- 5 and 6 were placed for sternal closing in Group A. In Group B, retrosternal sternal band at intercostal spaces 3 and five standard 5/0 steel wires at intercostal spaces 1- 2- 4-5 and 6 were placed. Then cutaneous and subcutaneous tissues were closed standardly. Following post-operative hemodynamic stability, all patients were ex-tubed at hour 4 to 6 and pain index scoring was obtained following mobilization during post-operative intensive unit care. Pain index was obtained by using pain scoring method Visual Analogue Scale (VAS) to evaluate the effect of both bands on post-operative pain. Patients were verbally asked to

rate the presence, frequency and severity of pain on the sternum by a scale of 1 to 10. Pain index scoring was calculated by multiplying post-operative pain frequency by pain severity.

Statistical analysis was performed by SPSS 15.0 version for Windows (SPSS Inc., Chicago, Illinois, USA) software. T-Test, Pearson Chi-Square test and Mann-Whitney test were used for data analysis. Statistical data was expressed as mean  $\pm$  standard deviation and significance was considered as  $p < 0.05$ .

The study was approved by Süleyman Demirel University Faculty of Medicine Scientific Study Projects Consulting Coordinating Board by project number B.30.2.SDU.0.20.05.07-507/5077 on 27.12.2011 and board decision no. 27/6.

**Table 2.**

	Group A	Group B
On pump CABG	45	52
Valve + CABG	3	2
Off- pump CABG	6	3
<i>CABG: Coronary artery bypass graft</i>		

## Results

Pre-operative demographics were not statistically significantly different between two groups (**Table 1**). Surgery type of operated patients is shown on **Table 2**. When peri- and post-operative variables were evaluated, there was no statistical difference in respect of cardiopulmonary bypass time (CPB), aortic cross clamp time (ACC), DR drainage amount, the amount of blood utilized, administration of inotropes, drain removal

**Table 3.**

	Group A ( n=54)	Group B ( n= 60)	P
CPB	100.1 $\pm$ 26	103.5 $\pm$ 30.2	NS
ACC	60.8 $\pm$ 10.2	63.6 $\pm$ 7.0	NS
Total drain (ml)	850 $\pm$ 330.5	800 $\pm$ 310.2	NS
Drain removal (hour)	44 $\pm$ 4.1	43 $\pm$ 4.08	NS
Superficial wound infection	-	-	-
Mediastinitis	-	-	-
Dehiscence	2	1	NS
Discharge time (days)	7	7	NS
Pain index	6 ( min 1- max 48)	4.00 (min.1 – max 63)	p=0.011, P <0.05
<i>Pain index: Pain severity x Pain frequency</i>			

time, superficial wound infection, mediastinitis, dehiscence and discharge time (  $p > 0.05$ ), (**Table 3**).

Mean pain index was 6.00 (min 1 – max 48 ) and 4.00 (min 1 – max 63) in Group A and Group B, respectively. Statistically significant difference was found between two groups in respect of pain index ( $p = 0.011$ ).

## Discussion


Complete closing of sternum following median sternotomy during open cardiac surgery, is a critical point in prevention of postoperative complications due to sternotomy. These complications include mainly dehiscence as well as a broad range of conditions such as pain, superficial wound infection and mediastinitis. Currently several sternal closing methods were described to prevent these complications.<sup>(4)</sup> Therefore, in our study we compared two different sternal bands used for sternal closing.

In patients with sternotomy, sternal dehiscence is the most frequent complication and the rate is 0.3% to 5%. This is a mechanical problem associated with certain factors such as lack of attention of patient to protect the sternum post-operatively, COPD, obesity, osteoporosis, use of bilateral mammarian arteries, renal insufficiency, administration of steroids and re-operation.<sup>(4)</sup> The incidence of sternal wound infection is 0.4-5% and it is closely associated with dehiscence.<sup>(10)</sup>


Closing of sternum is an important factor for sternal dehiscence. During sternal closing done by conventional approach using steel wires 5/0, bone cortex erosion, fracture and consequently sternal dehiscence can occur due to compression of steel wires. Sternal dehiscence may lead to respiratory dysfunction, infection, increased pain and re-exploration.<sup>(16)</sup> Use of sternal band, especially retrosternal application at intercostal space 3 and 4 to transversely surrounding whole sternum, is the most important mechanism to reduce development of sternal dehiscence with its 6 times larger surface than steel wires and thus providing equal loading on sternum. In our study, sternal dehiscence and sternal wound infection were not detected in both groups.

Postoperative pain is an acute pain gradually reducing by tissue healing accompanied also by inflammatory process.<sup>(6)</sup> Indeed, acute pain is a complex sensation besides its simple perceptual character. Difference in pain perception may be due to central process, fear, anxiety, depression and previous experience on pain perception. Certain patients may not describe the pain due to their higher pain tolerance or their introvert pain handling methods. Difference in pain perception of patients may be also due to post-operative analgesic administration methods. However, 30% of surgery patients don't require postoperative analgesics.<sup>(7)</sup> In a prospective clinical study, localization, distribution

**Figure 1.**




**Figure 2.**



and severity of postoperative pain were evaluated in consecutive 200 patients with cardiac surgery by median sternotomy.<sup>(8)</sup> Investigators determined the highest pain intensity on day 1 and lowest intensity on day 3. Although pain distribution was not different during post-operative period, there was difference in its localization.<sup>(9,15)</sup> Another factor affecting the pain severity is the age of patient. Young patients reported more severe

**Figure 3.**



pain than patients over 60 years.<sup>(8,11)</sup>

Previous studies comparing sternal band and conventional sternal closing also showed that usage of sternal band was associated with decreased pain and this was explained by reduced development of sternal dehiscence. However, in our study, we determined a difference between different methods of sternal bands in respect of pain development. When two groups of our study were compared, there was significant reduction in post-operative pain development in Group B compared to Group A. At this point, we consider that reduced compression of band method on a smaller bone cortex area only at intercostal space 3 in Group B compared to band method of Group A, could explain lower pain.

However, we think that the limited number of studies on pain in cardiac surgery and multifactorial nature of pain are limitations of our study. Although our study is prospective and including low number of patients, we conclude that larger studies will provide more guidance by including psychological and anatomic-physical parameters such as difference in physical load on bone structure by the material utilized.

## References

1. A.Rahman,O Burma.The effect of coronary artery bypass grafting with Cardiopulmonary bypass and beating heart technique on cardiac performance Turkish J Thorac Cardiovasc Surg 2001;9:68-73
2. H.Ekin,V.Kutay Tek ve İki Damar Hastalarında Çalışan Kalpte ve Kardi-yopulmoner Bypass ile Yapılan Myokardiyal Revaskülarizasyonların Karşılaştırılması Van Tıp Dergisi: 2004;11 (3):92-97
3. Cameron A, Davis KB, Green G, Schaff HV. Coronary bypass surgery with internal-thoracic-artery grafts-effects on survival over a 15-year period.N Engl J Med 1996;334:216-9.
4. Friedrich-Christian Riess,1 Nizar Awwad,1 Bettina Hoffmann. A Steel Band in Addition to 8 Wire Cerclages Reduces the Risk of Sternal Dehis-cence after Median Sternotomy. The Heart Surgery Forum 7 (6), 2004 [Epub September 2004]
5. Wewers ME, Lowe NK. A critical review of visual analogue scales in the measurement of clinical phenomena. Research in Nursing & Health 1990;13: 227-236,
6. Uyar M. Postoperatif ağrılı hastanın değerlendirilmesi ve ağrı ölçümü. Yücel A ed. Postoperatif analjezi 1. basım. İstanbul: Mavimer Matbacılık Yayıncılık Ltd. Şti. 2004; 27-36.
7. Rawal N. Postoperatif Ağrının Psikolojik Etkileri Editör: Erdine S. Ağrı. Nobel Tıp Kitabevleri 1.baskı. 2000; 124-125.
8. Mueller XM, Tinguely F, Tevaearai HT ve ark. Pain location, distribu-tion and intensity after cardiac surgery. Chest 2000; 118: 391-6.
9. Moore R, Follette DM, Berkoff HA. Poststernotomy fractures and pain management in open cardiac surgery. Chest 1994; 106: 1339-42.
10. Blanchard A, Hurni M, Ruchat P, et al: Incidence of deep and superficial sternal infection after open heart surgery: A ten years retrospective study from 1981 to 1991. Eur J Cardiothorac Surg 1995; 9: 153-8.
11. Farinas MC, Peralta FG, Bernal JM, et al: Suppurative mediastinitis after open heart surgery: A case-control study covering a seven year period in Santander, Spain. Clin Infect Dis 1995; 20: 272-6.
12. He GW, Ryan WH, Acuff TE, et al: Risk factors for operative mortality and sternal wound infection in bilateral internal mammary artery graft-ing. J Thorac Cardiovasc Surg 1994; 107: 196-9.
13. Kluytmans JAJW, Mouton JW, Ijzermann EPF, et al: Nasal carriage of staphylococcus aureus as a major risk factor for wound infections after cardiac surgery. J Infect Disc 1995; 171: 216-9.
14. Jurkiewicz MJ, Bostwick J, Hester TR, Bishop JB, Craver J. Infected median sternotomy wound. Ann. Surg. 1980; 191(6):738-43.
15. Catarino PA, Chamberlain MH, Wright NC, Black E. High-Pressure suc-tion drainage via a polyurethane foam in the management of postster-notomy mediastinitis. Ann. Thorac Surg. 2000; 70: 1891-5.
16. Robisceck F, Fokin A, Cook J, Bhatia D. Sternal Instability After Midline Sternotomy. Thorac Cardiovasc Surg. 2000; 48 (4): 1-8.

Received: 16/05/2017

Accepted: 04/09/2017

Published: 15/09/2017

### Disclosure and conflicts of interest:

The authors declare no conflict of interest.

### Corresponding author:

Kenan Abdurrahman Kara

Mail: kenankaradoc@gmail.com

*Kara K.A., Erk Z., Koçyiğit A., Gülmen Ş., Öcal A., Okutan H. Comparison between two sternum closing techniques by median sternotomy during open cardiac surgery EJCM 2017; 05 (3): 44-49. Doi: 10.15511/ejcm.17.00344.*

# Hybrid treatment of aortic bifurcation aneurysm

Funda Yıldırım<sup>1</sup>, Dilşad Amanvermez Şenarslan<sup>1</sup>, Adnan Taner Kurdal<sup>2</sup>,  
Barış Bayram<sup>1</sup>, Tülün Öztürk<sup>3</sup>, Mustafa Cerrahoğlu<sup>4</sup>, İhsan İşkesen<sup>4</sup>, Ömer Tetik<sup>4</sup>

<sup>1</sup>) Manisa Celal Bayar University Hafsa Sultan Hospital Cardiovascular Surgery Department, MD., Manisa, Turkey

<sup>2</sup>) Manisa Celal Bayar University Hafsa Sultan Hospital Cardiovascular Surgery Department, Assist.Prof., Manisa, Turkey

<sup>3</sup>) Manisa Celal Bayar University Hafsa Sultan Hospital Department of Anesthesiology and Reanimation, Prof.Dr., Manisa, Turkey

<sup>4</sup>) Manisa Celal Bayar University Hafsa Sultan Hospital Cardiovascular Surgery Department, Prof.Dr., Manisa, Turkey

## Abstract

The success of endovascular treatments mostly depends on anatomical suitability. Secure neck and iliac landing zones are required to avoid of endoleaks. Common iliac artery aneurysms frequently extend to the iliac bifurcation and complicates endovascular interventions. We report a hybrid operation of an abdominal bifurcation and extremely angulated iliac artery aneurysm that aneurysmatic segments involving both internal iliac arteries (IIA). We excluded the aneurysm by an uniiliac endovascular stent graft and contralateral IIA flow was supplied by retrograde filling from femoro-femoral bypass after ligation contralateral common iliac artery via retroperitoneal approach. It is important to consider alternative treatment modalities and in cases with extreme anatomic variations, to choose a hybrid endovascular approach may be the most feasible and less invasive.

**Keywords:** Endovascular treatments, hybrid operation, abdominal bifurcation, iliac artery aneurysm.

*Yıldırım F., Amanvermez Şenarslan D., Kurdal A. T., Bayram B., Öztürk T., Cerrahoğlu M., İşkesen İ., Tetik Ö Hybrid treatment of aortic bifurcation aneurysm EJCM 2017; 05 (3): 50-53. Doi: 10.15511/ejcm.17.00350.*

## Introduction

Infrarenal abdominal aortic aneurysm and iliac artery aneurysm are seen mostly together. The success of endovascular treatments mostly depends on anatomical suitability. Secure neck and iliac landing zones are required to avoid of endoleaks. Common iliac artery aneurysms (CIA) frequently extend to the iliac bifurcation and complicate the endovascular interventions. The preservation of circulation of at least one internal iliac artery (IIA) is highly recommended.

## Case


We report a hybrid operation of an abdominal bifurcation and extremely angulated iliac artery aneurysm that aneurysmatic segments involves both internal iliac arteries. A 71-year old man was diagnosed incidentally aortoiliac aneurysm by routine urologic examination and referred to our clinic (**Figure 1**). We planned hybrid approach. Both common femoral arteries were explored and snared. The aneurysm is excluded by a aorto-uniiliac endovascular stent graft system (Endurant II, Medtronic™) that extending to the right external iliac artery (EIA) and right IIA is closed with the extension graft. A femoro-femoral bypass is made with

8 mm ringed PTFE graft to supply left lower extremity circulation (**Figure 2, 3**). Then we explored the left common aneurysmatic iliac artery by retroperitoneal approach with lateral oblique abdominal incision. We ligated the CIA with nylon tape to prevent the enlargement of aneurysmal sac of the left CIA. The flow to the left IIA was supplied by retrograde filling from femoro-femoral bypass. Then we observed the decreased tension of the aneurysmal sac in left CIA also the left IIA pulse was palpated. No proximal, distal, or graft junction endoleaks occurred. The patient was discharged without trouble at postoperative fourth day.

## Discussion

EVAR has become the firstline choice in abdominal aortic aneurysm repair. But there are some limitations to its use; hostile neck anatomy, hostile iliac anatomy (HIA) and aneurysmal enlargement involving bilateral internal iliac artery. Approximately 30% of patients treated by EVAR have ectatic or aneurysmal common iliac arteries not suitable for distal sealing zones. Bilateral internal iliac artery occlusion during endovascular repair may lead significant morbidity, including gluteal claudication, erectile dysfunction, ischemia of the sigmoid colon and perineum and spinal cord injury.<sup>(1,2,3)</sup>

**Figure 1.** Computed tomography angiography 3 dimensional reconstruction view, aortic aneurysm just prior to aortic bifurcation and extending to common iliac artery. Aneurysmatic segment involves bilateral internal iliac artery and hostile iliac arteries.



In CIA aneurysms involving the IIA, there are some ways of preserving flow to IIA; Surgical revascularization of IIA via retroperitoneal or retroinguinal incision. An uncovered stent can extend the sealing zones, whilst maintaining complete preservation of pelvic circulation and offers support to the covered stent-graft.<sup>(4)</sup> Iliac side-branched graft (IBG) or sandwich technique can maintain pelvic perfusion. However, these adjunctive procedures increase the risk of endoleaks and they need to be applied by an experienced interventionist because of its technical difficulties.<sup>(5-8)</sup>

The presence of narrow iliac arteries (diameter <7 mm), hemodynamic iliac stenosis or obstruction, severe or extensive calcifications (>50%iliac circumferential calcification), angulations (>90 degree), previous aortailiac/femoral graft was defined as HIA.<sup>(5)</sup> HIA or previous surgical or endovascular graft greatly limits the endograft's maneuverability.<sup>(9)</sup>

This case have an aortic aneurysm just prior to aortic bifurcation and extending to CIA. The case is an



example to hostile iliac artery because of severe iliac angulation and also it is complicated by aneurysmatic segment involving bilateral IIA ostium. The aneurysm is treated in combined fasion: endovascular and conventional surgery. We used this approach for two reasons. First, to prevent recurrent king or stenosis of the left iliac endograft system due to extreme angulation to the right side. Secondly, to preserve the blood supply of at least one of the internal iliac artery.

We think that it is important to consider alternative treatment modalities and in cases with extreme anatomic variations, to choose a hybrid endovascular approach may be the most feasible and less invasive.

### Conclusion

Despite progress in endovascular therapy, large infrarenal to use combined thoracic and abdominal stent graft as an alternative treatment modality when the patient is not suitable for standard EVAR and open surgery.

**Figure 2, 3.** Insertion of uniiliac stent graft system.





## References

1. Faries PL, Morrissey N, Burks JA, Gravereaux E, Kerstein MD, Teodorescu VJ et al. Internal iliac artery revascularization as an adjunct to endovascular repair of aortoiliac aneurysms. *J Vasc Surg* 2001; 34(5): 892-9.
2. Cochenec F, Marzelle J, Allaire E, Desgranges P, Becquemin JP. Open vs endovascular repair of abdominal aortic aneurysm involving the iliac bifurcation. *J Vasc Surg* 2010; 51(6): 1360-6.
3. Bosanquet DC, Wilcox C, Whitehurst L, Cox A, Williams IM, Twine CP; British Society of Endovascular therapy (BSET). Systematic Review and Meta-analysis of the Effect of Internal Iliac Artery Exclusion for Patients Undergoing EVAR. *Eur J Vasc Endovasc Surg* 2017; 53(4): 534-548.
4. Silingardi R, Tasselli S, Gennai S, Saitta G, Coppi G. Endovascular preservation of pelvic circulation with external iliac-to-internal iliac artery "cross-stenting" in patients with aorto-iliac aneurysms: a case report and literature review. *J Cardiovasc Surg (Torino)* 2012; 53(5): 651-5.
5. Zeng Q, Huang L, Huang X, Peng M. Endovascular repair of abdominal aortic aneurysm with severely angulated neck and tortuous artery access: case report and literature review. *BMC Surg* 2015; 15: 20.
6. Noel-Lamy M, Jaskolka J, Lindsay TF, Oreopoulos GD, Tan KT. Internal Iliac Aneurysm Repair Outcomes Using a Modification of the Iliac Branch Graft. *Eur J Vasc Endovasc Surg* 2015; 50(4): 474-9.
7. Lim CS, Naji Y, Hussain ST, Pleban E, Wiszniewski A, Onida S et al. Modified Sandwich-graft Technique Employing Aorfix and Viabahn Stent-grafts to Preserve Hypogastric Flow in Cases of Complex Aortoiliac and Isolated Common Iliac Artery Aneurysms Including the Internal Iliac Artery Ostium. *Eur J Vasc Endovasc Surg* 2016; 51(3): 364-70.
8. Donas KP, Torsello G, Pitoulias GA, Austermann M, Papadimitriou DK. Surgical versus endovascular repair by iliac branch device of aneurysms involving the iliac bifurcation. *J Vasc Surg* 2011; 53(5): 1223-9.
9. Gallitto E, Gargiulo M, Faggioli G, Pini R, Mascoli C, Freyrie A et al. Impact of iliac artery anatomy on the outcome of fenestrated and branched endovascular aortic repair. *J Vasc Surg*. 2017; 6. pii: S0741-5214(17)31359-9. doi: 10.1016/j.jvs.2017.04.063. [Epub ahead of print] PubMed PMID: 28888759.

Received: 16/01/2017

Accepted: 22/07/2017

Published: 15/09/2017

### Disclosure and conflicts of interest:

The authors declare no conflict of interest.

### Corresponding author:

MD. Funda Yıldırım

**Mail:** fundanizamoglu@yahoo.com

*Yıldırım F., Amanvermez Şenarslan D., Kurdal A. T., Bayram B., Öztürk T., Cerrahoğlu M., İşkesen İ., Tetik Ö Hybrid treatment of aortic bifurcation aneurysm EJCM 2017; 05 (3): 50-53. Doi: 10.15511/ejcm.17.00350.*

# Extra-anatomic left carotico-subclavian artery bypass technique in a patient with subclavian steal syndrome: Case report

Ertan Demirdaş<sup>1</sup>, Kıvanç Atılğan<sup>2</sup>, Zafer Cengiz Er<sup>3</sup>, Ferit Çiçekçioğlu<sup>1</sup>

<sup>1)</sup> Bozok University Medicine Faculty Cardiovascular Surgery Department, Prof.Dr., Yozgat-Turkey

<sup>2)</sup> Bozok University Medicine Faculty Cardiovascular Surgery Department, Assist.Dr., Yozgat-Turkey

<sup>3)</sup> Bozok University Medicine Faculty Cardiovascular Surgery Department, Op. Dr., Yozgat-Turkey

## Abstract

Subclavian steal syndrome (SSS) is characterized by total occlusion of proximal part of subclavian artery resulting with retrograde flow of vertebral artery. The main risk factors in etiology are hypercholesterolemia and atherosclerosis. Congenital left subclavian artery stenosis and patent ductus arteriosus accompanying aortic coarctation, posttraumatic injuries, thromboembolism and arteritis are some of the rare factors causing SSS. SSS is usually an asymptomatic disease. However, the major symptoms are dizziness, claudication intermittent of upper extremity, headache and paresthesia in distal part of upper extremities. We are presenting a 57-year-old man presenting with a claudication intermittent on the left arm, paresthesia in fingers, dizziness and headache for a three year time. After applying a computerized tomographic angiography, we observed that the left subclavian artery was totally occluded at the origin point. There are several surgical approaches for the treatment of this syndrome. In this case, we preferred the carotico-subclavian bypass technique with supraclavicular approach due to its low rates of mortality and morbidity, and high rates of long term durability.

**Key words:** Graft, carotid artery, subclavian steal, supraclavicular

*Demirdaş E., Atılğan K., Er Z. C., Çiçekçioğlu F. Extra-anatomic Left Carotico-Subclavian Artery Bypass Technique in a patient with Subclavian Steal Syndrome: Case report EJCM 2017; 05 (3): 54-57. Doi: 10.15511/ejcm.17.00354.*

## Introduction

Subclavian steal syndrome (SSS) is usually an asymptomatic disease characterized by total occlusion etiology of proximal part of subclavian artery.<sup>(1)</sup> SSS was first described in the early 20th century and many of researches have been made both in diagnosing and treating areas. It is well known that SSS is a consequence of retrograde flow of vertebral artery following the total occlusion of proximal part of subclavian artery.<sup>(2)</sup> The main risk factors in etiology are hypercholesterolemia and atherosclerosis.<sup>(2-4)</sup> It is also possible to observe SSS due to congenital left subclavian artery stenosis and patent ductus arteriosus accompanying aortic coarctation.<sup>(5,6)</sup> Besides, posttraumatic injuries, thromboembolism and arteritis are some of the rare factors causing SSS.<sup>(1-3)</sup>

SSS is usually an asymptomatic disease. However, the major symptoms are dizziness, claudication intermittent of upper extremity, headache and paresthesia in distal part of upper extremities. SSS is often diagnosed incidentally by means of observing the retrograde flow of vertebral artery in colored doppler ultrasonography of carotid and vertebrobasillary arteries getting applied to the patients having dizziness and observing a gradient in tension arterial (TA) pressures between two upper extremities.

**Image 1. Preoperative Computerized Tomographic Angiography image**




## Case Report

A 57-year-old male with a history of claudication intermittent on left arm, paresthesia in dita, disseness and headache for three years applied to our department. Distal pulses were not palpable on the left upper extremity and TA was 130/80 mmHg on right arm and 60/30 mmHg on left arm. On the computerized tomographic angiography (CTA), we observed that the first 3 cm part of the subclavian artery beginning from the aortic arcus origin was totally occluded.**(Image 1)**

## Operation Technique

Invasive arterial monitorization was applied from right upper extremity. The patient was positioned as the head in hyperextension and the face looking to the opposite side, left shoulder was elevated and the upper extremities were pulled down. An 8 cm left supraclavicular incision was applied, common carotid and subclavian arteries were dissected gently and the proximal and distal parts were fixed with tapes. After systemic heparinization common carotid artery was clamped. Following arteriotomy the proximal segment of an 8 mm PTFE graft was anastomosed to the artery. After declamping, subclavian artery was clamped and arteriotomy was applied. Distal segment of PTFE graft was anastomosed to the subclavian artery.**(Image 2)** After

**Image 2. Intraoperative image of the graft**



the operation on both sides of upper extremities distal pulses were palpable. The patient was discharged on the postoperative 6th day without any complication.

## Discussion

Subclavian steal syndrome (SSS) is characterized by total occlusion of proximal part of subclavian artery resulting with retrograde flow of vertebral artery. The retrograde vertebral artery flow may cause variable symptoms like cerebellar ischaemia (dizziness, blumia), claudication intermittent and paresthaesia on the upper extremity.

There are different types of surgical techniques, almost all of which get a remarkable success in eliminating the symptoms, and surgery should be a treatment approach only in symptomatic patients. The first successful transthoracic intervention for a subclavian artery obstruction was achieved by De Bakey et al in 1958.<sup>(7)</sup> In 1964 Parrot described the carotico-subclavian artery bypass procedure.<sup>(8)</sup> Carotico-subclavian bypass, subclavio-subclavian bypass, axillo-subclavian bypass and subclavio-carotid transposition techniques are described as extra-anatomic procedures, and aorta-subclavian bypass and endarterectomy are described as anatomic procedures. Transthoracic incision techniques are not preferred anymore due to high mortality and morbidity rates.<sup>(9)</sup>

Percutaneous transluminal angioplasty and stenting

(PTAS) are some of the treatment approaches as an alternative to surgery. Especially PTAS have become the first option in SSS cases occurring as a result of vasculitis or radiotherapy.<sup>(10)</sup> However, the long term durability of surgery is superior to PTAS.<sup>(11)</sup>

Carotico-subclavian artery bypass technique is an extra-anatomic bypass process and has a low morbidity and mortality rates. Vogt et al, reported the mortality rate of intrathoracic techniques as 15% and of extra-anatomic techniques as 0%, in 1982.<sup>(12)</sup> Saphenous and artificial grafts can be used for bypassing. However, 5 year durability rates are reported as 95,2% for PTFE grafts, 83,9% for Dacron grafts and 64% for saphenous vein grafts.<sup>(13)</sup> In our case we preferred an 8 mm PTFE graft.

In recent researches, carotico-subclavian artery bypass technique is described as a process having low mortality and morbidity, and long term survival rates.<sup>(14,15)</sup> Normal carotid artery anatomy is a necessity for this technique, otherwise it is possible to cause a carotid artery steal syndrome.<sup>(16-18)</sup>

## Conclusion

Extra-anatomic carotico-subclavian artery bypass technique is a relatively easy process in comparison to other techniques, and has better mortality and morbidity rates, because of that we prefer and recommend this technique as a first option for patients having SSS.

## References

1. Potter BJ, Pinto DS. Subclavian steal syndrome. Circulation, 2014 Jun 3; 129(22):2320-3
2. Abova V, Kamineni A, Allison Ma, Mc Dermott MM, Crouse JR, Ni h, Szklo et al. The epidemiology of subclavian stenosis and its association with markers of subclinical atherosclerosis: The Multi-Ethnic study of atherosclerosis (MESA). Atherosclerosis. 2010;211:266-70.
3. Labropoulos N, Nandivada P, Bekelis K. Prevalence and impact of the subclavian steal syndrome. Ann Surg. 2010;252:166-70.
4. Harper C, Cardullo PA, Weyman AK, Patterson RB. Transcranial Doppler ultrasonography of the basilar artery in patients with retrograde vertebral artery flow. J Vasc Surg. 2008;48:859-64.
5. Gerber N. Congenital atresia of the subclavian artery producing an subclavian steal syndrome. Amer j dis child 1967;113:709-13.
6. Daves j, Treger A. Vertebral grand larcency. Circulation 1964;29:91-3.
7. DeBakey ME, Morris GC, Jordan GL, Cooley DA. Segmental thrombo-obliterative disease of branches of aortic arch. JAMA 1958;166:988-1003.
8. Parrot JD. The subclavian steal sendrome. Arch Surg 1964;88:661-5.
9. Fields WS, Lemak HA. Joint study of extracranial Arterial Occlusion, VII. Subclavian steal - a review of 168 cases. JAMA 1972;222:1139-43.
10. Kim HJ, Lee CS, Kim JS, Know SU, Kim JL, Park JW, et al. Outcomes after endovascular treatment of symptomatic patients with Takayasu's arteritis. Interv Neuroradiol 2011;17:252-60.
11. Tetik Ö, YAŞA H, Demir T, Gürbüz A. Subklavyan Steal sendromu: 3 olgu nedeniyle. İzmir Atatürk Eğitim ve araştırma Hastanesi Tıp dergisi 2004;139-41
12. Vogt DP, Hertzner NR, O'Hara PJ, Beven EG. Brachiocephalic arterial reconstruction. Ann surg. 1982;196:541-52.
13. Law MM, Colburn MD, Moore WS, Quinones-Baldrich WJ, Machleder HI, Gelabert HA. Carotid-subclavian bypass for brachiocephalic occlusive disease. Stroke 1995;26:1565-71.
14. Aburahma AF, Robinson PA, Jennings TG. Carotid-subclavian bypass grafting with polytetrafluoroethylene grafts for syptomatic subclavian artery stenosis or occlusion: A 20-year experience. J vasc surg 2000;32:411-9.
15. Vitti MJ, Thompson BW, Read RC, et al. Carotid-subclavian bypass: A twenty-two year experience. J Vasc surg 1994;20:411-8.
16. Lord RS, Ehrenfeld WK. Carotid-subclavian bypass: A hemodynamic study. Surgery 1969;66:521-6.
17. Cook CH, Stemmer EA, Connolly JE. Effect of peripheral resistance on carotid blood flow after Carotid-subclavian bypass. Arch surg 1972;105:9-13.
18. Swayngim DM Jr, Avellone JC. Hemodynamic consequences of axillo-axillary bypass J cardiovasc surg 1982;23:65-8.

Received: 09/02/2017

Accepted: 31/08/2017

Published: 15/09/2017

### Disclosure and conflicts of interest:

The authors declare no conflict of interest.

### Corresponding author:

Prof. Dr. Ertan Demirdaş

**Mail:** dr.ertandemirdas@gmail.com

*Demirdaş E., Atılgan K., Er Z. C., Çiçekçiöğlü F. Extra-anatomic Left Carotico-Subclavian Artery Bypass Technique in a patient with Subclavian Steal Syndrome: Case report EJCM 2017; 05 (3): 54-57. Doi: 10.15511/ejcm.17.00354.*

# Use of new oral anticoagulants secondary to Paget–Schroetter syndrome in a young sportsman: A case report

Kenan Abdurrahman Kara<sup>1</sup>, Uğur Diliçıkık<sup>2</sup>, Başar Sarıkaya<sup>3</sup>, Şenol Gülmen<sup>4</sup>

<sup>1)</sup> Hisar Intercontinental Hospital, Cardiovascular Surgery Department, Op.Dr., Istanbul, Turkey

<sup>2)</sup> Kanuni Sultan Süleyman Training and Research Hospital, Department of Sports Medicine, MD., Istanbul, Turkey

<sup>3)</sup> Yeditepe University, Department of Radiology, Assoc.Prof., Istanbul, Turkey

<sup>4)</sup> Süleyman Demirel University Faculty of Medicine, Department of Cardiovascular Surgery, Assoc. Prof.Dr., Isparta, Turkey

## Introduction

**Introduction:** The Paget–Schroetter syndrome or “effort thrombosis” is a deep venous thrombosis of subclavian-axillary vein at the costoclavicular junction . It affects predominately young and athletic people with an inherent anatomic variant at the costoclavicular junction that triggers the formation of a primary thrombosis. It is important to identify this syndrome in order to prevent further complications, such as post-thrombotic syndrome, pulmonary embolism (PE), or even death. We present a clinical case in which a young professional basketball player developed a subclavian venous thrombosis and was managed with new oral anticoagulant.

**Keyword:** Oral anticoagulants, effort thrombosis, Paget–Schroetter syndrome, sportsman

*Kara K.A., Diliçıkık U., Sarıkaya B., Gülmen Ş Treatment of Paget-Schroetter syndrome with a new anticoagulant therapy in young sportsman: A case report. EJCM 2017; 05 (3): 58-61. Doi: 10.15511/ejcm.17.00358.*

## Case Report

We describe a case of a 17-year-old male, Turkish basketball player, who presented to the clinic with one-week course of left upper limb pain with swelling and mild erythema. There was no previous history of chest trauma and patient denied having fever. The patient had no known medical problems, no medications and no prior family history of hematologic-related diseases. On physical examination his blood pressure was 116/88 mmHg, pulse rate 88/min, pulse oximetry showed 94% on room air, temperature was 36.4°C. His lungs were clear on auscultation and his heart sounds were normal. Edema, tenderness and erythema were found over the left upper extremity, peripheral arterial pulses were clear with palpation and no motor or sensitive deficit was present. The patient had also Urschel's sign, characterized by the dilation of veins that can be visible across the shoulder and upper arm.<sup>(1,2)</sup>


An upper extremity venous Doppler ultrasound study was performed which confirmed the presence of a partially occlusive deep vein thrombosis in the left

subclavian vein. Hematology Investigations, including complete blood count, coagulation studies, and renal function tests, yielded normal results. Work-up for thrombophilic disorders included homocysteine, protein C, antithrombin III, and anti-cardiolipin antibody levels which were all within the normal range. In the ED, the patient was then placed on Pradaxa (dabigatran) 150mg orally daily

## Discussion

Primary axillary-subclavian vein thrombosis, first described by Paget in 1875 and Schroetter in 1884, was named Paget-Schroetter syndrome (PSS) by Hughes in 1948.<sup>(7)</sup> Upper extremity effort thrombosis accounts for approximately 1–4% of all episodes of venous thrombosis.<sup>(2)</sup> PSS most often develops among young adults who work in occupations that require repeated arm movements which cause axillo-subclavian vein trauma and facilitate the development of deep vein thrombosis (DVT).<sup>(8)</sup> Several sporting activities have been associated with axillary vein thrombosis, including ball games, racket games, and aquatic sports. Due to repeti-

**Figure 1.**



tive extension and hyperabduction movements of the upper limb, this kind of thrombosis is known as effort thrombosis.<sup>(9)</sup> Hematologic, anatomic, and iatrogenic predisposing factors such as thrombophilia, thoracic outlet syndrome (TOS), and major vein catheterisation are typically underlying conditions.<sup>(10-13)</sup> Clinical features of PSS include sudden onset of pain, swelling, edema, and cyanosis of the upper limb. Collateral veins are evident around the shoulder and chest. The pain increases with exercise and decreases with rest and elevation of the affected extremity.<sup>(14)</sup>

Two theories explain the initial process that generates the final event of thrombosis in the Paget-Schroetter syndrome. The first one has been involved in a frequent activity of the arm followed by the development of a hypertrophy in the anterior scalene muscle; this, leading to blood stasis and clot formation.

Secondly, several descriptions suggest that secondary thrombosis begins with a structural compression (abnormalities in the thoracic outlet), which triggers a continuous process of fibrosis and scarring in the external environment surrounding the subclavian vein, of an exchange of the collagen fibers from loose connective tissue to a dense collagen scar. These perpetuate the process of stasis and thrombosis.<sup>(3)</sup>

The typical clinical presentation is a sudden onset pain, heaviness, blue-red discoloration and swelling of the upper affected arm. However, due to the mainly presentation in young athletic patients, the syndrome tends to be asymptomatic, or described by the patient as a simple muscular strain. In patients with intermittent obstruction, the symptoms will appear and disappear according to the process of obstruction. Occasionally, patients present the “Urschel’s sign”, characterized by the dilation of veins that can be visible across the shoulder and upper arm.<sup>(2)</sup>

The mainstream treatment of upper extremity throm-

bosis is based on the correction of the underlying defect and pre-vents future episodes. Currently, the gold standard in the management of this unusual thrombosis is the thrombolytic therapy. Catheter-directed thrombolysis has reported a successful of 62–82%, being higher in recent fresh clots (symptoms <2–6 weeks) followed by a venoplasty if there is evidence of residual obstruction.<sup>(2,4)</sup> Anticoagulation therapy reduces the overall mortality and morbidity associated with this syndrome. However, some studies demonstrate an increased risk of residual venous obstruction in 78% of cases managed only with anticoagulation.<sup>(4)</sup> Urschel et al. describes that only 29% of patients treated with anticoagulation reported a good or excellent outcome.<sup>(5)</sup> There was no evidence related to the use of new oral anticoagulants for the management of upper limb thrombosis.

In cases of external compression by scarring and fibrosis the treatment needs a surgical decompression to avoid a subsequently rethrombosis.<sup>(6)</sup> The surgical procedure should be performing 1 or 2 month after the episode.

## Conclusion

In conclusion, early diagnosis and treatment of the Paget-Schroetter syndrome is critical for preventing potentially fatal complications such as pulmonary embolism. Prophylaxis is important for preventing recurrent thrombosis and for avoiding development of post-thrombotic syndrome. Paget-Schroetter syndrome should be considered a possible cause of painful swelling of the upper limb, especially significant edema, blue discoloration, and dilatation of the superficial veins in the patient’s right upper limb in young, active patients who use their arms excessively.

In addition, there are few studies related with the use of new oral anticoagulants for medical management. In this case report, we use one of the new oral anticoagulants. Further clinical trials are needed to show the clinical benefits of this drug and prove an association.



## References

1. Illig KA, Doyle AJ. A comprehensive review of Paget-Schroetter syndrome. *J Vasc Surg* 2010 Jun;51(6):1538–47.
2. Alla VM, Natarajan N, Kaushik M, Warriar R, Nair CK. Paget-schroetter syndrome: review of pathogenesis and treatment of effort thrombosis. *West J Emerg Med* 2010 Sep;11(4):358–62.
3. Zell L, Kindermann W, Marschall F, Scheffler P, Gross J, Buchter A. Paget-Schroetter syndrome in sports activities--case study and literature review. *Angiology* 2001 May;52(5):337–42.
4. Adams JT, DeWeese JA. "Effort" thrombosis of the axillary and subclavian veins. *J Trauma* 1971 Nov;11(11):923–30.
5. Urschel HC Jr, Razzuk MA. Paget-Schroetter syndrome: what is the best management? *Ann Thorac Surg* 2000 Jun;69(6):1663–8; discussion 1668–9.
6. Beygui RE, Olcott C 4th, Dalman RL. Subclavian vein thrombosis: outcome analysis based on etiology and modality of treatment. *Ann Vasc Surg* 1997 May;11(3):247–55.
7. Oktar GL, Ergul EG: Paget-Schroetter syndrome. *Hong Kong Med J*, 2007,13: 243–245.
8. Ilhan E, Ture M, Yilmaz C, et al.: Subclavian vein thrombosis extending into the internal jugular vein: Paget-von Schroetter Syndrome. *J Clin Med Res*, 2009, 1: 178–180.
9. Maharaj D, Ramdass M, Perry A, et al.: Paget von Schroetter Syndrome secondary to exotic dancing: a case study. *Int J Angiol*, 2003, 12: 143–144.
10. Akita K, Ibukuro K, Yamaguchi K, et al.: The subclavius posticus muscle: a factor in arterial, venous or brachial plexus compression? *Surg Radiol Anat*, 2000, 22: 111–115.
11. Ozçakar L, Kaymak B, Turan S, et al.: Thoracic outlet syndrome, Paget-Schroetter syndrome and aberrant subclavian artery in a young man. *Joint Bone Spine*, 2006, 73: 469–471.
12. Ozçakar L, Dönmez G, Yörübulut M, et al.: Paget-Schroetter syndrome forerunning the diagnoses of thoracic outlet syndrome and thrombophilia. *Clin Appl Thromb Hemost*, 2010, 16: 351–355.
13. Lewandowski A, Syska-Sumińska J, Dłużniewski M: [Pulmonary embolism suspicion in a young female patient with the Paget-von Schrötter syndrome]. *Kardiol Pol*, 2008, 66: 969–971.
14. Bolcal C, Oz K, Sirin G, et al.: Paget-von Schroetter Syndrome associated with thrombophilia. *Turk J Thorac Cardiovasc Surg*, 2007, 15: 304–306.

Received: 16/06/2017

Accepted: 04/09/2017

Published: 15/09/2017

### Disclosure and conflicts of interest:

The authors declare no conflict of interest.

### Corresponding author:

Kenan Abdurrahman Kara

**Mail:** kenankaradoc@gmail.com

*Kara K.A., Diliçik U., Sarıkaya B., Gülmen Ş Treatment of Paget-Schroetter syndrome with a new anticoagulant therapy in young sportsman: A case report. EJCM 2017; 05 (3): 58-61. Doi: 10.15511/ejcm.17.00358.*

# Endovascular treatment of a ruptured abdominal aortic aneurysm in a patient with cardiogenic shock

Ali Baran Budak<sup>1</sup>, Atike Tekeli Kunt<sup>2</sup>, Orhan Eren Günertem<sup>2</sup>, Naim Boran Tümer<sup>2</sup>, Kanat Özışık<sup>3</sup>, Serdar Günaydın<sup>3</sup>

<sup>1)</sup> Ankara Numune Education and Research Hospital, Cardiovascular Surgery Department, Exp. Dr., Ankara, Turkey

<sup>2)</sup> Ankara Numune Education and Research Hospital, Cardiovascular Surgery Department, Assist. Dr., Ankara, Turkey

<sup>3)</sup> Ankara Numune Education and Research Hospital, Cardiovascular Surgery Department, Prof. Dr., Ankara, Turkey

## Abstract

Abdominal aort aneurysm is an asymptomatic and potentially lethal disease which its rupture is an emergent situation. In many centers, patients presenting with ruptured abdominal aortic aneurysm (rAAA) are most often treated with open repair (OR). A 55-year-old male with acute abdominal pain admitted to our hospital. Computed tomography scan showed a rAAA. A sudden cardiac arrest occurred. The patient immediately transferred to catheter laboratory with cardiogenic shock table and endovascular treatment was performed. Then he was discharged in the postoperative 7th day.

**Keyword:** ruptured abdominal aort aneurysm, endovascular repair, cardiogenic shock

*Budak A. B., Tekeli Kunt A., Günertem O. E., Tümer N. B., Özışık K., Günaydın S. Endovascular treatment of a ruptured abdominal aortic aneurysm in a patient with cardiogenic shock EJCM 2017; 05 (3): 62-66. Doi: 10.15511/ejcm.17.00362.*

## Introduction

Abdominal aortic aneurysm is an asymptomatic and potentially lethal disease which its rupture is an emergent situation and has a fatality rate of 100% if untreated. Misdiagnosis causes delays for the treatment and it can be catastrophic. Depending on the patient's status during admission, the mortality ranges from 20% to 70% even in treated patients.<sup>[1,2]</sup> In many centers, patients presenting with ruptured abdominal aortic aneurysm (rAAA) are most often treated with open repair(OR). But in centers whom have a hybrid operating room as the surgeons' comfort and confidence with the technique increases, ruptured endovascular aneurysm repair (rEVAR) has become the preferred treatment option. Furthermore, patients with rAAA who are transferred for a higher level of care have superior clinical outcomes when treated with rEVAR, compared with OR.<sup>[3]</sup> Herein, we aimed to demonstrate the feasibility of emergent rEVAR in a patient with cardiogenic shock who underwent successful resuscitations after three times cardiac arrest.

## Case Report

A 55-year-old male with acute abdominal pain was transferred to our emergency department and took under evaluation of general surgery and urology clinics with suspicious of renal colic and ileus. At the present-


tation, serum urea level was 80 mg/dl, creatinin level was 2.97 mg/dl and hemoglobin level was 7.5 g/dl. Computed tomography scan showed a rAAA extending from the caudal of superior mesenteric artery (SMA) to aortic bifurcation with a diameter of 8.53 cm. The aneurysmal sac included the right renal artery, but left renal artery could not be demonstrated (**Figure 1**). The measurements of the aneurysm is shown in **Table 1**. The level of the rupture is shown in **Figure 2**.

After computed tomographic assessment, a sudden cardiac arrest occurred. The patient was entubated and an immediate resuscitation procedure was performed. Systolic blood pressure could be increased to 40 mmHg and cardiac arrest repeated for 3 times. After the systolic blood pressure was secured around 50 mmHg, the patient immediately transferred to catheter laboratory for endovascular repair. Bilateral surgical cutdown to the groin was performed. after positioning a 20 F sheath

**Table 1.** The measurements of the aneurysm.

Measured Area	Millimeter
Proximal diameter of neck	30.3
Distal diameter of neck	28
Neck length	12
Max aneurysm diameter	85.3
Distal aorta diameter	22.5
SMA to distal aorta	142.4
Proximal diameter of RCI	13
Diameter of RCI	14.5
Distal diameter of RCI	18.2
Length of RCI	73.2
Total length from right (SMA to right internal iliac artery)	215.7
Proximal diameter of LCI	12
LCI diameter	13.5
Distal diameter of LCI	16.9
Length of LCI	76.8
Total length from left (Infra renal level to left internal iliac artery)	219.2

**Figure 1.** 3D CT Image. Aneurysm extends from the distal of SMA and includes the renal arteries



to left CFA (Sentrart Sheath, Medtronic), a uniiliac stent graft sized 32x14x102 mm was positioned (Endurant® II uniiliac, Medtronic). SMA was preserved. After successful positioning of the graft, a 16x20x156 mm stent graft (Endurant® II, Medtronic) was used for the extension to the left iliac artery. Particular attention was paid to keep the left internal iliac artery open. During the procedure, cannulation of the contralateral limb was unable to be achieved because of the tortuous aortoiliac course. Since the patient was unstable, the Endurant system was combined with Talent® Converter stent-graft occluder (20mm, Medtronic, Inc.; Minneapolis, Minn) positioned to right common iliac artery via a 18.5 F introducer sheath. An aortography was made and a minimal leak from the overlap area was observed. Upon this, Reliant® (Medtronic, Inc.; Minneapolis, Minn) balloon catheter was used to expand the stent graft and reinforce the overlap zone. After the closure of the rAAA was confirmed. (Figure 3) Additional left-to-right femorofemoral bypass with a 8mm Gore-Tex graft was performed. After successful closure of the rAAA, the patient was taken to the intensive care unit. Postprocedural hemoglobin was 9.8 g/dl, serum creatinin level was 3.63 mg/dl. The patient was extubated 32 h after the procedure without neuro-

logic dysfunction. Renal failure was expected and the patient was consulted to the nephrology clinic for routine dialysis program. The patient was discharged in the postoperative 7th day.

## Discussion


rAAA is a catastrophic condition and the role of rEVAR is still controversial due to the absence of well-supported evidences. Mortality is higher after OR than EVAR according to retrospective studies<sup>[4-7]</sup> and confirmed by the Swedish Vascular Registry.<sup>[8]</sup> Furthermore, preoperative shock table and unstable status was found as a major factor predicting mortality rate.<sup>[9]</sup> Also many publications pointed out that the duration of intensive care is shorter<sup>[10,11]</sup> and operative blood loss and/or need for transfusion is lower in rEVAR patients.<sup>[12,13]</sup> Our patient was followed up for 5 days in ICU and we only gave 2 units of erythrocyte suspension and 2 units of fresh frozen plasma during the operation. Since rEVAR allows the reduction of intraoperative blood loss and the elimination of the aortic clamping and fluid shifts/losses due to a large transperitoneal incision, we did not encounter with a cardiac complication. After overlapping the stent grafts, we observed a type III

**Figure 2.** Computed-Tomography Scan. Level of rupture was seen.



endoleak from the overlap zone then we fixed it by using a balloon catheter. We also placed an occluder to right common iliac artery to prevent a type II endoleak, as mentioned above. Immediate resuscitation in the emergency department, successful management of a comatose and unstable patient both in the preoperative

**Figure 3. Angiographic Image. Successful closure of the rAAA was seen.**



and postoperative period protected the patient to have a neurological complication.

Our first goal was to achieve a safe transfer to the intensive care unit and keep patient alive. Due to lack of every size and types of graft in our hospital and inability to perform a contralateral limb cannulation we had to admit renal complications and ischemia risk of contralateral limb. Femorofemoral bypass surgery prevented ischemic complications but renal failure was occurred and the patient became dialysis dependent. On the other hand renal artery occlusion is also a well known complication of EVAR and it can occur by several mechanism.<sup>[14-16]</sup>

In this case report we aimed to demonstrate the feasibility of emergent rEVAR in a patient with cardiogenic shock and help cardiovascular surgeons for the future. In a this kind of emergent status choosing the treatment modality (open surgery whether endovascular repair) is debatable. Our another goal was to enrich the literature about this kind of medical scenarios. Further studies and long-term results of endovascular procedures in patients with ruptured aort aneursym are still needed.

## References

1. Larsson E, Granath F, Swedenborg J et al (2008) More patients are treated for nonruptured abdominal aortic aneurysms, but the proportion of women remains unchanged. *J Vasc Surg* 48(4):802–807
2. Ozdemir BA, Karthikesalingam A, Sinha S et al (2015) Association of hospital structures with mortality from ruptured abdominal aortic aneurysm. *Br J Surg* 102(5):516–524
3. Mandawat A, Mandawat A, Sosa JA, Muhs BE, Indes JE. Endovascular repair is associated with superior clinical outcomes in patients transferred for treatment of ruptured abdominal aortic aneurysms. *J Endovasc Ther* 2012;19(1):88–95.
4. Lo RC, Bensley RP, Hamdan AD et al (2013) Gender differences in abdominal aortic aneurysm presentation, repair, and mortality in the Vascular Study Group of New England. *J Vasc Surg* 57(5): 1261–1268
5. Mehta M, Byrne WJ, Robinson H et al (2012) Women derive less benefit from elective endovascular aneurysm repair than men. *J Vasc Surg* 55(4):906–913
6. de Leur K, Flu HC, Ho GH et al (2015) Outcome of elective treatment of abdominal aortic aneurysm in elderly patients. *Int J Surg* 15:117–123
7. Ullery BW, Tran K, Chandra V et al (2015) Association of an endovascular-first protocol for ruptured abdominal aortic aneurysms with survival and discharge disposition. *JAMA Surg* 150(11):1058–1065
8. Swedvasc (2015) Swedvasc. Annual report from the Swedish National vascular registry. In: SSVS U (ed.) <http://www.ucr.uu.se/swedvasc/>
9. Mehta M, Byrne J, Darling RC 3rd, Paty PS, Roddy SP, Kreinberg PB, et al. Endovascular repair of ruptured infrarenal abdominal aortic aneurysm is associated with lower 30-day mortality and better 5-year survival rates than open surgical repair. *J Vasc Surg* 2013;57(2):368–75.
10. Wibmer A, Schoder M, Wolff KS et al: Improved survival after abdominal aortic aneurysm rupture by offering both open and endovascular repair. *Arch Surg*, 2008; 143(6): 544–49; discussion 550
11. Anain PM, Anain JM Sr., Tiso M et al: Early and mid-term results of ruptured abdominal aortic aneurysms in the endovascular era in a community hospital. *J Vasc Surg*, 2007; 46(5): 898–905
12. Anain PM, Anain JM Sr., Tiso M et al: Early and mid-term results of ruptured abdominal aortic aneurysms in the endovascular era in a community hospital. *J Vasc Surg*, 2007; 46(5): 898–905
13. Starnes BW, Quiroga E, Hutter C et al: Management of ruptured abdominal aortic aneurysm in the endovascular era. *J Vasc Surg*, 2010; 51(1): 9–17; discussion 18
14. Gorich J, Rilinger N, Soldner J, et al. Endovascular repair of aortic aneurysms: treatment of complications. *J Endovasc Surg* 1999;6:136–14
15. Liewald F, Sharrer-Pamler R, Gorich J, et al. Intraoperative, perioperative, and late complications with endovascular therapy of aortic aneurysm. *Eur J Vasc Endovasc Surg* 2001; 22:251–256
16. Hausegger KA, Schedlbauer P, Deutschmann HA, Tiesenhäusen K. Complications in endoluminal repair of abdominal aortic aneurysms. *Eur J Radiol* 2001;39:22–33

Received: 30/06/2017

Accepted: 26/08/2017

Published: 15/09/2017

### Disclosure and conflicts of interest:

The authors declare no conflict of interest.

### Corresponding author:

Dr. Ali Baran Budak

**Mail:** [drbaranbudak@gmail.com](mailto:drbaranbudak@gmail.com)

E-Journal of Cardiovascular Medicine welcomes scientific contributions in the field of cardiovascular and thoracic surgery - all aspects of surgery of the heart, vessels and the chest in various article types: new ideas, brief communications, work in progress, follow-up studies, original articles, best evidence topics, case reports, reports on unexpected results etc. All manuscripts shall be reviewed by the Editor-in-Chief, Associate Editors, Invited Referees and a Statistician when appropriate. If accepted, articles will be posted online and opened up for discussion. Acceptance criteria are based on the originality, significance, and validity of the material presented.

All material to be considered for publication in E-Journal of Cardiovascular Medicine should be submitted in electronic form via the journal's online submission system. (<http://my.ejmanager.com/ejcm/>)

A cover letter should be enclosed to all new manuscripts (to be filled in online), specifying the name of the journal and the type of paper, and including the following statements:

- The manuscript should not be previously published in print or electronic form and is not under consideration by another publication.
- All authors should contribute to the content of the article.
- All authors should read and approve the submission of the manuscript to ICVTS.
- Subject to acceptance, authors will sign an exclusive license to publish.
- No ethical problem or conflict of interest should exist.

If your first language is not English, we recommend that you consult an English language editing service to ensure that the academic content of your paper is fully understood by journal editors and reviewers. Language editing does not guarantee that your manuscript will be accepted for publication.

Manuscripts should be prepared using a word-processing package (save in .doc, .docx or .rtf format). The font type and font size should preferably be Arial or Times New Roman 11 points. The manuscript should be double-spaced and should include line and page numbers. The lines of the reference list do not need to be numbered; include a section break before.

### Manuscripts should be organized as follows:

(a) Title page; (b) Abstract and Key words; (c) Text with the following sections (not applicable for article types with unstructured abstracts): Introduction, Materials and Methods, Results, Discussion, Acknowledgement (optional), Funding statement, Conflict of interest statement, (d) Figure (and Video) legends; (e) Tables; (f) References.

**Title page (1st page):** Title: should be brief and descriptive (100 characters) - no abbreviations are allowed, even if well known.

**Authors:** list all authors by full first name, initial of or full middle name and family name. Qualifications are not required. Ensure the author names correspond (in spelling and order of appearance) with the metadata of the system

**Institution(s):** include the name of all institutions with the location

(department, institution, city, country) to which the work should be attributed (in English). Use superscript numbers to connect authors and their department or institution.

**Corresponding author:** The full name, full postal address, telephone/fax numbers and the e-mail address should be typed at the bottom of the title page.

**Meeting presentation:** If the manuscript was (or will be) presented at a meeting, include the meeting name, venue, and the date on which it was (or will be) read; also indicate if you have submitted an Abstract of this manuscript for the EACTS or ESTS annual meeting and whether it has been accepted (if known).

**Word count:** The total number of words of the whole article (including title page, abstract, main text, legends, tables and references) must be specified on the title page.

**Abstract (2nd page):** An abstract should be a concise summary of the manuscript. Reference citations are not allowed. The abstract should be factual and free of abbreviations, except for SI units of measurement.

**Keywords:** Following the abstract, 3-6 keywords should be given for subject indexing.

**Introduction:** Should state the purpose of the investigation and give a short review of pertinent literature.

**Materials and methods:** Should be described in detail with appropriate information about patients or experimental animals. Use of abbreviations renders the text difficult to read; abbreviations should be limited to SI units of measurement and to those most commonly used, e.g. VSD, ASD, CABG (abbreviations should not be included in headings and extensions should be included at first mention).

**Results:** Results should be reported concisely and regarded as an important part of the manuscript. They should be presented either in tables and figures, and briefly commented on in the text, or in the text alone. Repetition of results should be avoided!

**Discussion:** The discussion is an interpretation of the results and their significance with reference to pertinent work by other authors. It should be clear and concise.

**Acknowledgement:** Acknowledgements and details of non-financial support must be included at the end of the text before the references. Personal acknowledgements should precede those of institutions or agencies.

**Tables:** All tables must be included in the manuscript file, should start on separate pages and be accompanied by a title, and footnotes where necessary. The tables should be numbered consecutively using Arabic numerals. Units in which results are expressed should be given in parentheses at the top of each column and not repeated in each line of the table.

**References:** Authors are responsible for checking the accuracy of all references. If you use EndNote or Reference Manager to facilitate referencing citations (not required for submission), this journal's style is available for use. References should be numbered in order of appearance in the text (in Arabic numerals in parentheses) and must be listed numerically in the reference list. Journal titles and author initials should be properly abbreviated and punctuated.

## GENERAL RULES

Files should be prepared as a Word document using font size 12 Times New Roman characters, double-spaced and with 2.5 cm margins on each side, top and bottom. Only standard abbreviations should be used; other shortened phrases should be indicated in parentheses as used in the text. Generic or chemical names of drugs should be used instead of trade names.

## ETHICAL ISSUES

### Publishing responsibilities of authors and Ethics

The publication of an article in a peer-reviewed journal is an essential building block in the development of a coherent and respected network of knowledge. It is a direct reflection of the quality of work of the author and the institutions that support them. Peer-reviewed articles support and embody the scientific method. It is therefore important to agree upon standards of expected ethical behavior.

### Reporting standards

Authors of reports of original research should present an accurate account of the work performed as well as an objective discussion of its significance. Underlying data should be represented accurately in the paper. A paper should contain sufficient detail and references to permit others to replicate the work. Fraudulent or knowingly inaccurate statements constitute unethical behavior and are unacceptable. Review and professional publication articles should also be accurate and objective, and editorial 'opinion' works should be identified as such.

### Hazards and human or animal subjects

If the work involves chemicals, procedures or equipment that have any unusual hazards inherent in their use, the author must clearly identify these in the manuscript. If the work involves the use of animal or human subjects, the author should ensure that the manuscript contains a statement that all procedures were performed in compliance with relevant laws and institutional guidelines and that the appropriate institutional committee(s) has approved them. Authors should include a statement in the manuscript that informed consent was obtained for experimentation with human subjects. The privacy rights of human subjects must always be observed.

### Use of patient images or case details

Studies on patients or volunteers require ethics committee approval and informed consent, which should be documented in the paper. Appropriate consents, permissions and releases must be obtained where an author wishes to include case details or other personal information or images of patients and any other individuals in publication. Written consents must be retained by the author and copies of the consents or evidence that such consents have been obtained must be provided to us on request. Particular care should be taken with obtaining consent where children are concerned (in particular where a child has special needs or learning disabilities), where an individual's head or face appears, or where reference is made to an individual's name or other personal details.

## Originality and plagiarism

The authors should ensure that they have written entirely original works, and if the authors have used the work and/or words of

others, that this has been appropriately cited or quoted. Plagiarism takes many forms, from 'passing off' another's paper as the author's own paper, to copying or paraphrasing substantial parts of another's paper (without attribution), to claiming results from research conducted by others. Plagiarism in all its forms constitutes unethical publishing behavior and is unacceptable.

## Data access and retention

Authors may be asked to provide the raw data in connection with a paper for editorial review, and should be prepared to provide public access to such data (consistent with the ALPSP-STM Statement on Data and Databases), if practicable, and should in any event be prepared to retain such data for a reasonable time after publication.

## Multiple, redundant or concurrent publication

An author should not in general publish manuscripts describing essentially the same research in more than one journal or primary publication. Submitting the same manuscript to more than one journal concurrently constitutes unethical publishing behavior and is unacceptable. In general, an author should not submit for consideration in another journal a previously published paper. Publication of some kinds of articles (e.g. clinical guidelines, translations) in more than one journal is sometimes justifiable, provided certain conditions are met. The authors and editors of the journals concerned must agree to the secondary publication, which must reflect the same data and interpretation of the primary document. The primary reference must be cited in the secondary publication.

## Acknowledgement of sources

Proper acknowledgment of the work of others must always be given. Authors should cite publications that have been influential in determining the nature of the reported work. Information obtained privately, as in conversation, correspondence, or discussion with third parties, must not be used or reported without explicit, written permission from the source. Information obtained in the course of confidential services, such as refereeing manuscripts or grant applications, must not be used without the explicit written permission of the author of the work involved in those services.

## Fundamental errors in published works

When an author discovers a significant error or inaccuracy in his/her own published work, it is the author's obligation to promptly notify the journal editor or publisher and cooperate with the editor to retract or correct the paper. If the editor or the publisher learns from a third party that a published work contains a significant error, it is the obligation of the author to promptly retract or correct the paper or provide evidence to the editor of the correctness of the original paper.

## Authorship of the paper

Authorship should be limited to those who have made a significant contribution to the conception, design, execution, or interpretation of the reported study. All those who have made significant



contributions should be listed as co-authors. Where there are others who have participated in certain substantive aspects of the research project, they should be acknowledged or listed as contributors. The corresponding author should ensure that all appropriate co-authors and no inappropriate co-authors are included on the paper, and that all co-authors have seen and approved the final version of the paper and have agreed to its submission for publication.

### Changes to authorship

This policy concerns the addition, deletion, or rearrangement of author names in the authorship of accepted manuscripts. Before the accepted manuscript is published in an online issue:

Requests to add or remove an author, or to rearrange the author names, must be sent to the Journal Manager by the corresponding author of the accepted manuscript, and must include:

The reason the name should be added or removed, or the author names rearranged

Written confirmation (e-mail, fax, letter) from all authors that they agree with the addition, removal or rearrangement. In the case of addition or removal of authors, this includes confirmation from the author being added or removed

Requests that are not sent by the corresponding author will be forwarded by the Journal Manager to the corresponding author, who must follow the procedure described above.

#### Note that:

- Journal Managers will inform the Journal Editors of any such requests
- Publication of the accepted manuscript in an online issue is suspended until authorship has been agreed
- After the accepted manuscript has been published in an online issue:

Any requests to add, delete or rearrange author names in an article published in an online issue will follow the same policies as noted above and may result in a corrigendum.

## TYPES OF PAPERS

### Original Articles

Original articles should consist of sections titled as “Abstract, Introduction, Materials and Methods, Results, Discussion and Conclusion”. For information about the abstract, refer to ‘Manuscript Formatting’ section.

The Introduction section of the manuscript should clearly state the purpose of the manuscript and include a brief summary of the most relevant national and international literature stating the main purposes and research question of the study. Contradictory aspects of the research, if present, should be mentioned. The expected contribution of this study to family medicine and practice should be highlighted.

The Materials and Methods section should describe the study population and the study design, with adequate information on the

techniques, materials and methods used. The section should include information of the study type, population, sample, sample size and selection of the sample. Validity and reliability of scales and questionnaires used also should be referred to. A clear description of the statistical methods should also be given.

The Results section should include a detailed report on the findings of the study. All figures, tables and illustrations should be used in this section. Results should be presented either as text or figures and/or tables and not be replicated.

The Discussion section of the study should emphasize the importance of the results and compare them with the results of other authors with relevant citations from the most recent literature. Study limitations and strengths should be specified. Suggestions for further studies in this area should be added.

The Conclusion should include the main conclusions based on the results of the research, emphasize the contributions of the study to family practice and propose original suggestions. A brief revision of all the results and the discussion should be avoided.

Original articles excluding case reports and systematic reviews should not exceed 3000 words excluding the abstract, references and tables. Case reports should not exceed 1000 words excluding the abstract, references and tables. There are no restrictions for systematic reviews.

### Short Reports

Short Reports are accepted when the research topic, aim and results of the study are limited in scope and in cases that do not require writing a full original article. Short Reports can be described as a summarized version that have been prepared according to the structure of research articles. Publishing an article as a short report does not reflect a lower quality. The same rules as relevant to original articles apply to preparing a short report, but structured abstracts are not mandatory references and tables should not exceed 6 and 2 in number, respectively. Abstracts should not exceed 100 words and the text should be restricted to a maximum of 1000 words.

### Reviews

Reviews are evidence-based articles about a specific topic using relevant citations from the most recent literature with the authors’ conclusions on this subject. The author is expected to have conducted research on the subject and to have experience in order to discuss and analyze the subject. There is no obligation to follow a particular format and may contain subtitles depending on the subject. The text should not exceed 4000 words excluding the title, abstracts, references and tables. E Journal of Cardiovascular Medicine, only publishes review articles solicited by the editors.

### Letters to Editor and Comments

Letters to the editor or comments can be sent to provide commentary and analysis concerning an article published in the journal, to give information about ongoing research, to provide informa-

tion in cardiology and cardiovascular-vascular-endovascular surgery, cardio-metabolic and vascular sciences. Letters to the editor or comments may include an optional title, tables and references. These articles should not exceed 1000 words.

## What Would You Do?

These are brief articles discussing cases and situations encountered in cardiology and cardiovascular surgery with a biopsychosocial approach. If necessary, photographs (with permission from the patient/owner) may be added. Sections should consist of a title, case report, discussion, questions and answers. Brief comments can be sent to provide commentary on previous articles and case reports written by other authors. Comments should include the number of the journal the article was published in. The text should not exceed 1000 words.

## International Reprints

Translations of important documents, declarations and guidelines prepared by international organizations in the field of cardiology and cardiovascular surgery, may be published in the journal. Presubmission Inquiry to the Editorial Board of the Journal before submitting the article is recommended. It is the translator's responsibility to obtain permission from the owner of the original manuscript for publication and translation.

## News

These articles focus on advances and innovations in clinical topics relevant to cardiology and cardiovascular surgery. There is no obligation to follow a particular format. The text should be limited to 1000 words.

## Editorials

Editorials usually provide information about the editorial policy of E Journal of Cardiovascular Medicine, give commentary and feedback on articles published in the journal, draw attention to topics of current interest and give information related to and discuss the development of cardiology and cardiovascular surgery in the world. They are mainly written by the members of the Editorial Board. Editorials are limited to 2000 words with some exceptions and may include a title and references when necessary.

## MANUSCRIPT FORMATTING

Manuscripts should be designed in the following order:

*Title page*

*Abstract*

*Main text*

*References*

*Tables, figures and illustrations*

### Title Page

The title page of the manuscript should include: The title, first

and last names of each author. Complete affiliation and title for each author, with the name of department (s) and institution (s) to which the work should be attributed.

The corresponding author should be clearly identified with name, address, telephone- facsimile number and email address for correspondence about the manuscript. Authors should clearly indicate if the article has previously been presented at a congress or scientific meeting. The title should be concise and informative without abbreviations and not exceed 10 words.

### Abstract

Abstracts should be exact in English, with a minimum of 150 and maximum of 350 words. Abstracts of original research articles should be structured under subheadings as follows: objectives, methods, results and conclusion. A maximum of 3 key words should be added to English abstracts.

### Text

The text contains the rest of the manuscript. It is structured differently according to the type of manuscript (original research article, review, etc.). For example, original research articles should consist of aim and objectives, methods, results, discussion and conclusion.

### References

References should be cited in consecutive numerical order as first mentioned in the text and designated by the reference number in parentheses. If the number of authors for the reference is more than 6 authors, list the first three authors and add "et al".

Journal names should be abbreviated as used in Index Medicus. References should be cited in the Vancouver style. For detailed information please visit the relevant link

### Examples:

**For research articles follow the example below:**

– Verschuren WM, Jacobs DR, Bloemberg BP, et al. Serum total cholesterol and long-term coronary heart disease mortality. JAMA 1995; 274(2): 131–6.

For book chapters follow the example below:

– Rakel RE. The family physician. In: Rakel RE, editor. Textbook of family practice. 5th ed. Philadelphia: W.B. Saunders; 1995. p. 3-19.

**For web pages follow the example below:**

– Guidance for clinicians. An International Benchmarking Study. <http://www.who.int/topics/surgery/> accessed: 29/09/2002.

### Tables and Figures

Legends should take place on the top of the page for tables, and bottom of the page for figures and placed on separate pages. Explain all nonstandard abbreviations in footnotes.



**E Journal of Cardiovascular Medicine**  
is a global e-journal targeting articles on:

- clinical cardiology,
- interventional cardiology,
  - arrhythmia,
- cardiovascular surgery,
- vascular & endovascular surgery,
- vascular biology